

ABSTRACT OF THE DISCLOSURE

Method of identification and quantitative analysis of alcohol(s) in a sample by mass spectrometry using stable isotope labeled internal standard is provided. Said internal standard is prepared by reaction of an authentic sample of said alcohol with a stable isotope labeled reagent, and is added to a sample containing said alcohol. Said alcohol in said sample is then quantitatively converted to a chemical compound of identical structure, except the stable isotope atoms, as that of said internal standard using a non-labeled reagent. Said sample is then extracted and the extract is analyzed by mass spectrometry. Identification and quantification of said alcohol are made from a plot of ion ratio of said converted alcohol to said internal standard versus alcohol concentration.

REFERENCES

US patent documents

- | | | |
|-----------|--------------------|----------------------|
| 5,559,038 | September 24, 1996 | J. Fred Kolhouse |
| 6,358,996 | March 19, 2002 | Michael S. Alexander |

Other references

Mark M. Kushnir et al, "Comparison of four derivatizing reagents for 6-acetylmorphine GC-MS analysis", Journal of Analytical Toxicology, July/August 1999, page 262-269, vol. 23.

Michael L. Smith et al, "Forensic drug testing for opiates. VI. Urine testing for hydromorphone, hydrocodone, oxymorphone, and oxycodone with commercial opiate immunoassay and GC-MS", Journal of Analytical Toxicology, Jan/Feb 1995, page 18-26, vol. 19.

J. G. Guillot et al, "Determination of heroin, 6-acetylmorphine, and morphine in biological fluids using their propionyl derivatives with ion trap GC-MS", Journal of Analytical Toxicology, Mar/April 1997, page 127-133, vol. 21.